



# Cardiopulmonary Considerations for High School Student-Athletes During the COVID-19 Pandemic: Update to the NFHS-AMSSM Guidance Statement

*Endorsed by the American Medical Society for Sports Medicine (AMSSM) and National Federation of State High School Associations (NFHS).*

Jonathan A. Drezner, MD,<sup>1</sup> William M. Heinz, MD,<sup>2</sup> Irfan M. Asif, MD,<sup>3</sup> Casey G. Batten, MD,<sup>4</sup> Karl B. Fields, MD,<sup>5</sup> Neha P. Raukar, MD,<sup>6</sup> Verle D. Valentine, MD,<sup>7</sup> Kevin D. Walter, MD,<sup>8</sup> and Aaron L. Baggish, MD<sup>9</sup>

<sup>1</sup>Department of Family Medicine and Center for Sports Cardiology, University of Washington

<sup>2</sup>National Federation of State High School Associations

<sup>3</sup>Department of Family and Community Medicine, University of Alabama Birmingham

<sup>4</sup>Kerlan-Jobe Orthopaedic Institute

<sup>5</sup>Cone Health Sports Medicine

<sup>6</sup>Department of Emergency Medicine, Mayo Clinic

<sup>7</sup>Sanford Orthopedics & Sports Medicine

<sup>8</sup>Departments of Orthopaedic Surgery & Pediatrics, Medical College of Wisconsin

<sup>9</sup>Massachusetts General Hospital, Division of Cardiology, Cardiovascular Performance Program

## Correspondence:

Jonathan A. Drezner, MD

Professor, Department of Family Medicine

Director, UW Medicine Center for Sports Cardiology

University of Washington, Box 354060

3800 Montlake Blvd NE

Seattle, WA 98195

206-598-3294

[jdrezner@uw.edu](mailto:jdrezner@uw.edu)

## **DISCLAIMER – NFHS Position Statements and Guidelines:**

The NFHS regularly distributes position statements and guidelines to promote public awareness of certain health and safety-related issues. Such information is neither exhaustive nor necessarily applicable to all circumstances or individuals, and is no substitute for consultation with appropriate health-care professionals. Statutes, codes or environmental conditions may be relevant. NFHS position statements or guidelines should be considered in conjunction with other pertinent materials when taking action or planning care. The NFHS reserves the right to rescind or modify any such document at any time.

Keywords: cardiac; coronavirus; COVID-19; prevention; screening; sports

Cardiac injury from SARS-CoV-2 infection among hospitalized patients was reported early in the COVID-19 pandemic.<sup>10</sup> Concern and uncertainty regarding the risk of cardiac sequelae in young athletes with SARS-CoV-2 infection led to the development of several consensus recommendations for the cardiac evaluation of athletes following SARS-CoV-2 infection.<sup>1,2,4-6,9,11</sup> These guidelines were based on expert opinion and emerging clinical experience but lacked scientific data.

Recent large cohort studies in athletes have demonstrated a low risk of cardiac involvement and have greatly informed the use of cardiac testing after SARS-CoV-2 infection.<sup>3,7,8</sup> The Outcome Registry for Cardiac Conditions in Athletes (ORCCA) reported a 0.7% (95% CI: 0.4, 1.1) overall prevalence of cardiac involvement in 3,018 collegiate athletes from 42 universities that had largely undergone cardiac ‘triad’ testing with a resting 12-lead electrocardiogram (ECG), transthoracic echocardiogram (TTE), and troponin blood assay.<sup>8</sup> The ORCCA study also found that athletes with cardiopulmonary symptoms (e.g., chest pain, dyspnea, palpitations) during the acute illness or upon return to exercise were 3.1 (95% CI: 1.2, 7.7) times more likely to have cardiac involvement.<sup>8</sup> Similarly, a study of 789 professional athletes all of whom had undergone cardiac triad testing reported a 0.6% prevalence of cardiac inflammation.<sup>7</sup> Notably, all professional athletes diagnosed with myocarditis or pericarditis also had moderate symptoms defined as fever, flu-like illness, or cardiopulmonary symptoms.<sup>7</sup> Lastly, the Big Ten registry reported a 2.3% prevalence of clinical or subclinical myocardial involvement in 1,597 collegiate athletes who underwent mandatory screening cardiac magnetic resonance imaging (MRI) after SARS-CoV-2 infection.<sup>3</sup> Among the 13 universities participating, the reported prevalence ranged from 0 to 7.6%, with 3 sites reporting no myocardial involvement among 189 athletes.<sup>3</sup> This marked variation is likely not explained by the underlying pathological process but rather by technical and interpretation variability between sites and the relative absence of normative cardiac MRI data in young competitive athletes. Importantly, none of these large cohort studies in collegiate and professional athletes, despite on-going surveillance, have reported an adverse cardiac event associated with SARS-CoV-2 infection.

In light of these studies, an expert task force from the National Federation of State High School Associations (NFHS) and the American Medical Society for Sports Medicine (AMSSM) reconvened to update guidelines for the cardiac assessment of high school student-athletes with prior SARS-CoV-2 infection before sports participation (Figure 1).<sup>4</sup> In the absence of large cohort data in high school athletes, findings in college and professional athletes were extrapolated to the high school level. While many classifications of COVID-19 illness severity

have emerged, we used the definitions for mild, moderate, and cardiopulmonary symptoms as applied in large athlete cohort studies.<sup>3,7,8</sup> Key updates and recommendations include:

- *Asymptomatic and mild symptoms*: Athletes with asymptomatic infections or only mild symptoms (e.g., common cold-like symptoms without fever, gastrointestinal symptoms, or loss of taste/smell) do not require additional cardiac testing unless clinically indicated. Athletes should check-in with a clinician (e.g., physician, nurse practitioner, physician assistant, or athletic trainer) to determine if further clinical evaluation is needed. Athletes should be at least 3-5 days from symptom onset or positive test before beginning an exercise progression.
- *Moderate and cardiopulmonary symptoms*: Additional cardiac testing (e.g., ECG, TTE, troponin) should be considered in athletes with moderate symptoms (e.g., fever >100.4°F, chills, flu-like syndrome for ≥2 days) or initial cardiopulmonary symptoms (e.g., chest pain, dyspnea, palpitations). Athletes with remote infections and moderate symptoms >3 months ago who never received a work-up but have returned to full activity without symptoms do not need additional cardiac testing. Cardiology consultation and cardiac MRI should be considered for abnormal results and as clinically indicated. We recommend athletes are at least 5-7 days from symptom onset and that moderate symptoms are resolved before starting an exercise progression.
- *Cardiopulmonary symptoms on return to exercise*: All athletes with SARS-CoV-2 infections should be closely monitored for new cardiopulmonary symptoms as they return to exercise. Athletes with cardiopulmonary symptoms when they return to exercise (e.g., exertional chest pain, excessive dyspnea, syncope, palpitations, or unexplained exercise intolerance) should undergo additional cardiac testing (e.g., ECG, TTE, troponin) if not already performed and be evaluated by a cardiologist with consideration for a cardiac MRI or other investigations as indicated.
- *Return-to-sport exercise progression*: The return-to-sport progression and timeline should be individualized and is based on numerous factors including baseline fitness, severity and duration of COVID-19 symptoms, and tolerance to progressive levels of exertion. Most athletes will require a graded progression over at least a few days. Absent special indications, a prolonged return-to-sport timeline is not supported by evidence and further restriction from sports participation can contribute to detraining, increased injury risk, and mental health concerns.
- *Preparticipation Physical Evaluation (PPE)*: Additional history questions during a routine PPE should consider if the athlete had a COVID-19 illness. If yes, consider clarifying: when, what symptoms, and if the athlete is experiencing any new symptoms with exercise, especially chest pain.
- *Emergency Action Plan (EAP)*: The task force again stresses the importance of a well-rehearsed EAP for every sport at every venue with clear access to an Automated External Defibrillator (AED).

**Figure 1.** Cardiopulmonary considerations for high school student-athletes during the COVID-19 pandemic.

ECG, electrocardiogram; Echo, echocardiogram; GI, gastrointestinal; hs-Tn, high sensitivity troponin; MRI, magnetic resonance imaging


## References:

1. Baggish AL, Drezner JA, Kim JH, Martinez MW, Prutkin JM. The resurgence of sport in the wake of COVID-19: cardiac considerations in competitive athletes. *Br J Sports Med.* 2020;54:1130-1131.
2. Baggish AL, Levine BD. Icarus and sports after COVID 19: too close to the sun? *Circulation.* 2020;142:615-617.
3. Daniels CJ, Rajpal S, Greenshields JT, et al. Prevalence of Clinical and Subclinical Myocarditis in Competitive Athletes With Recent SARS-CoV-2 Infection: Results From the Big Ten COVID-19 Cardiac Registry. *JAMA Cardiol* [published online May 27, 2021]. doi: 10.1001/jamacardio.2021.2065
4. Drezner JA, Heinz WM, Asif IM, et al. Cardiopulmonary Considerations for High School Student-Athletes During the COVID-19 Pandemic: NFHS-AMSSM Guidance Statement. *Sports Health.* 2020;12:459-461.
5. Elliott N, Martin R, Heron N, Elliott J, Grimstead D, Biswas A. Infographic. Graduated return to play guidance following COVID-19 infection. *Br J Sports Med.* 2020;54:1174-1175.
6. Kim JH, Levine BD, Phelan D, et al. Coronavirus Disease 2019 and the Athletic Heart: Emerging Perspectives on Pathology, Risks, and Return to Play. *JAMA Cardiol.* 2021;6:219-227.
7. Martinez MW, Tucker AM, Bloom OJ, et al. Prevalence of Inflammatory Heart Disease Among Professional Athletes With Prior COVID-19 Infection Who Received Systematic Return-to-Play Cardiac Screening. *JAMA Cardiol.* 2021;6:745-752.
8. Moulson N, Petek BJ, Drezner JA, et al. SARS-CoV-2 Cardiac Involvement in Young Competitive Athletes. *Circulation* [published online April 17, 2021]. doi:10.1161/CIRCULATIONAHA.121.054824
9. Phelan D, Kim JH, Chung EH. A Game Plan for the Resumption of Sport and Exercise After Coronavirus Disease 2019 (COVID-19) Infection. *JAMA Cardiol.* 2020;5:1085-1086.
10. Shi S, Qin M, Shen B, et al. Association of cardiac injury with mortality in hospitalized patients with COVID-19 in Wuhan, China. *JAMA Cardiol.* 2020;5:802-810.
11. Wilson MG, Hull JH, Rogers J, et al. Cardiorespiratory considerations for return-to-play in elite athletes after COVID-19 infection: a practical guide for sport and exercise medicine physicians. *Br J Sports Med.* 2020;54:1157-1161.

**Update - August 2021**



# Cardiopulmonary Considerations for High School Student-Athletes during the COVID-19 Pandemic



- ECG should be compared to previous when available
- Troponin testing (hs-cTnI or cTnI) should be performed after 48 hours without exercise
- Confirmed myocarditis, pulmonary embolism, or other cardiopulmonary disorder should be managed per medical guidelines